


KEY ANSWERS

1	1	16	3	31	3	46	4
2	2	17	2	32	3	47	1
3	2	18	1	33	1	48	3
4	3	19	1	34	2	49	3
5	1	20	4	35	2	50	1
6	4	21	2	36	1	51	2
7	1	22	1	37	2	52	2
8	1	23	3	38	1	53	2
9	4	24	2	39	3	54	1
10	2	25	1	40	1	55	4
11	4	26	4	41	1	56	2
12	2	27	2	42	4	57	2
13	4	28	3	43	4	58	3
14	3	29	4	44	3	59	1
15	3	30	3	45	3	60	3


1. If r and r' denote the angles inside the prism having angle of prism 50° considering that during interval of time from $t = 0$ to $t = t$, r varies with time as $r = 10^\circ + t^2$. During this time r' will vary with time as



- 1) $40^\circ - t^2$ 2) $40^\circ + t^2$ 3) $50^\circ - t^2$ 4) $50^\circ + t^2$ **Ans.(1)**

Solution: $A = r + r'$, $r' = A - r = A - (10 + t^2) = A - 10 - t^2 = 50 - 10 - t^2 = 40^\circ - t^2$

2. If AB is incident plane wave front then refracted wave front is ($n_2 > n_1$)



- 1) 2) 3) 4) **Ans.(2)**

3. The total energy carried by the light wave when it travels from a rarer to a non-reflecting and non-absorbing medium

- 1) decreases 2) remains same 3) increases
3) either increases or decreases depending upon angle of incidence **Ans.(2)**

Solution: Remains same

4. If the radius of first Bohr orbit is r , then the radius of the second Bohr orbit will be

- (1) $2r$ 2) $8r$ 3) $4r$ 4) $2\sqrt{2}r$ **Ans.(3)**

Solution: $R_2 = n^2 r = 2^2 r = 4r$

5. Match the following types of nuclei with examples shown

	Column-I		Column-II
A	Isotopes	(i)	${}_3\text{Li}^7, {}_4\text{Be}^7$
B	Isobars	(ii)	${}_8\text{O}^{18}, {}_9\text{F}^{19}$
C	Isotones	(iii)	${}_1\text{H}^1, {}_1\text{H}^2$

- 1) A-iii, B-i, C-ii 2) A-ii, B-iii, C-i 3) A-i, B-iii, C-ii 4) A-iii, B-ii, C-i **Ans.(1)**

6. Which of the following statements is incorrect with reference to 'Nuclear force'?

- 1) Potential energy is minimum, if the separation between the nucleons is 0.8 fm
 2) Nuclear force becomes attractive for nucleon distances larger than 0.8 fm
 3) Nuclear force becomes repulsive for nucleon distances less than 0.8 fm
 4) Nuclear force is always attractive

Ans.(4)

7. The range of electrical conductivity (σ) and resistivity (ρ) for metals, among the following, is


- 1) $\rho \rightarrow 10^{-2} - 10^{-8} \Omega\text{m}$ 2) $\rho \rightarrow 10^{-5} - 10^6 \Omega\text{m}$ 3) $\rho \rightarrow 10^{11} - 10^{19} \Omega\text{m}$ 4) $\rho \rightarrow 10^2 - 10^8 \Omega\text{m}$
 $\sigma \rightarrow 10^2 - 10^8 \text{Sm}^{-1}$ $\sigma \rightarrow 10^5 - 10^{-6} \text{Sm}^{-1}$ $\sigma \rightarrow 10^{-11} - 10^{-19} \text{Sm}^{-1}$ $\sigma \rightarrow 10^{-2} - 10^{-8} \text{Sm}^{-1}$ **Ans.(1)**

8. Which of the following statements is correct for an n-type semiconductor?

- 1) The donor energy level lies just below the bottom of the conduction band
 2) The donor energy level lies closely above the top of the valence band
 3) The donor energy level lies at the half way mark of forbidden energy gap
 4) The donor energy level does not exist

Ans.(1)

9. The circuit shown in the figure contains two ideal diodes D_1 and D_2 . If a cell of emf 3 V and negligible internal resistance is connected as shown, then the current through 70Ω resistance (in ampere) is



- 1) 0 2) 0.01 3) 0.02 4) 0.03 **Ans.(4)**

Solution : D_1 is RB and D_2 is FB

$$I = \frac{V}{R_{\text{eff}} + r} = \frac{V}{30 + 70 + 0} = \frac{3}{100} = 0.03 \text{ A}$$

10. In determining the refractive index of a glass slab using a travelling microscope, the following readings are tabulated.


- (a) Reading of travelling microscope for ink mark = 5.123 cm
 (b) Reading of travelling microscope for ink mark through glass slab = 6.123 cm
 (c) Reading of travelling microscope for chalk dust on glass slab = 8.123 cm

From the data, the refractive index of a glass slab is

- 1) 1.390 2) 1.500 3) 1.601 (4) 1.399 **Ans.(2)**

$$\text{Solution : } n = \frac{R_3 - R_1}{R_3 - R_2} = \frac{8.123 - 5.123}{8.123 - 6.123} = \frac{3}{2} = 1.5$$

11. In an experiment to determine the figure of merit of a galvanometer by half deflection method, a student constructed the following circuit.



He unplugged a resistance of 52001Ω in R. When K_1 is closed and K_2 is open, the deflection observed in the galvanometer is 26 div. When K_2 is also closed and a resistance of 90Ω is removed in S, the deflection becomes 13 div. The resistance of galvanometer is nearly

- 1) 116.0Ω 2) 45.0Ω 3) 103.0Ω 4) 91.6Ω **Ans.(4)**

Solution: 91.6Ω (half deflection method).

12. While determining the coefficient of viscosity of the given liquid, a spherical steel ball sinks by a distance $h = 0.9$ m. The radius of the ball is $r = \sqrt{3} \times 10^3$ m. The time taken by the ball to sink in three trials are tabulated as follows.

Trial No.	Time taken by the ball to fall by h (in second)
1.	2.75
2.	2.65
3.	2.70

The difference between the densities of the steel ball and the liquid is 7000 kg m^{-3} . If $g = 10.0 \text{ ms}^{-2}$, then the coefficient of viscosity of the given liquid at room temperature is

- 1) 0.28 Pa.s 2) 0.14 Pa.s 3) $0.14 \times 10^{-3} \text{ Pa.s}$ 4) 14 Pa.s **Ans.(2)**

Solution : $v = \frac{2(\rho - \sigma)r^2g}{9\eta}$

$$\frac{0.9}{2.7} = \frac{2 \times 7000 \times 3 \times 10^{-6} \times 10}{9\eta}$$

$$\therefore \eta = 0.14 \text{ Pa s}$$

13. Which of the following expressions can be deduced on the basis of dimensional analysis? (All symbols have their usual meanings)

- (1) $s = ut + \frac{1}{2}at^2$ (2) $x = A \cos \omega t$ (3) $N = N_0 e^{-\lambda t}$ (4) $F = 6\pi\eta rv$ **Ans.(4)**

14. Two stones begin to fall from rest from the same height, with the second stone starting to fall ' Δt ' seconds after the first falls from rest. The distance of separation between the two stones becomes 'H'. ' t_0 ' seconds after the first stone starts its motion. Then t_0 is equal to

- 1) $\frac{H}{g\Delta t}$ 2) $\frac{H}{\Delta t} + \frac{\Delta t}{2g}$ 3) $\frac{H}{g\Delta t} - \frac{\Delta t}{2}$ 4) $\frac{H}{g\Delta t} + \frac{\Delta t}{2}$ **Ans.(3)**

Solution :

$$x_1 = \frac{1}{2}gt_0^2$$

$$x_2 = \frac{1}{2}g(t_0 + \Delta t)^2$$

$$x_2 - x_1 = H = \frac{1}{2}g(t_0 + \Delta t)^2 - \frac{1}{2}gt_0^2 \Rightarrow t_0 = \frac{H}{g\Delta t} - \frac{\Delta t}{2}$$

15. In the projectile motion of a particle on a level ground, which of the following remains constant with reference to time and position?

- 1) Vertical component of the velocity of the projectile
2) Average velocity between any two points on the path
3) Horizontal component of velocity
4) Angle between the instantaneous velocity with the horizontal

Ans.(3)

16. A particle is in uniform circular motion. The equation of its trajectory is given by $(x - 2)^2 + y^2 = 25$, where x and y are in meter. The speed of the particle is 2 ms^{-1} . When the particle attains the lowest 'y' co-ordinate, the acceleration of the particle is (in ms^{-1})

1) $0.4\hat{i}$ 2) $0.4\hat{j}$ 3) $0.8\hat{i}$ 4) $0.8\hat{j}$ **Ans. (3)**

Solution: $(x - 2)^2 + y^2 = 25$


Comparing with the equation of circle

$x^2 + y^2 = r^2$, we have $r = 5$

$$\therefore a = \frac{v^2}{r} = \frac{2^2}{5} = 0.8$$


At lowest point of y-coordinate, 'a' is along x-axis

17. A wooden block of mass M lies on a rough floor. Another wooden block of the same mass is hanging from the point O through strings as shown in the figure. To achieve equilibrium, the co-efficient of static friction between the block on the floor with the floor itself is



1) $\mu = \cos \theta$ 2) $\mu = \cot \theta$ 3) $\mu = \sin \theta$ 4) $\mu = \tan \theta$ **Ans.(2)**

Solution:




$$T \sin \theta = Mg \rightarrow (1)$$

$$T \cos \theta = \mu Mg \rightarrow (2)$$

$$\frac{(2)}{(1)} \cot \theta = \mu$$

18. A block of certain mass is placed on a rough floor. The coefficients of static and kinetic friction between the block and the floor are 0.4 and 0.25 respectively. A constant horizontal force $F = 20 \text{ N}$ acts on it so that the velocity of the block varies with time according to the following graph. The mass of the block is nearly (Take $g \approx 10 \text{ ms}^{-2}$)



1) 2.2 kg 2) 4.4 kg 3) 1.2 kg 4) 1.0 kg **Ans.(1)**

Solution : $\mu_s = 0.4$ $\mu_k = 0.25$

$$\text{Acceleration} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{20}{3}$$

$$F = ma + \mu mg$$

$$m = \frac{F}{\mu g + a} \approx 2.2 \text{ Kg}$$

19. A body of mass 0.25 kg travels along a straight line from $x = 0$ to $x = 2$ m with a speed $v = k x^{3/2}$ where $k = 2$ SI units. The work done by the net force during this displacement is
 (1) 4J (2) 8J (3) 16J (4) 32J **Ans.(1)**

Solution : $v = Kx^{\frac{3}{2}}$

$$w = \Delta KE = \frac{1}{2}mv^2 = \frac{1}{2}mk^2x^3 = \frac{1}{2} \times 0.25 \times 4 \times 2^3 = 4J$$

20. During an elastic collision between two bodies, which of the following statements are correct?
 I. The initial kinetic energy is equal to the final kinetic energy of the system.
 II. The linear momentum is conserved.
 III. The kinetic energy during Δt (the collision time) is not conserved.
 1) I and II only (2) II and III only (3) I and III only (4) I, II and III **Ans.(4)**

21. Three particles of mass 1kg, 2kg and 3kg are placed at the vertices A, B and C respectively of an equilateral triangle ABC of side 1 m. The centre of mass of the system from vertex A (located at origin) is


1) (0, 0) 2) $\left(\frac{7}{12}, \frac{3\sqrt{3}}{12}\right)$ 3) $\left(\frac{9}{12}, \frac{3\sqrt{3}}{12}\right)$ 4) $\left(\frac{7}{12}, \frac{6+3\sqrt{3}}{12}\right)$ **Ans.(2)**

Solution: $X_{cm} = \frac{m_1x_1 + m_2x_2 + m_3x_3}{m_1 + m_2 + m_3} = \frac{7}{12}m$

$$Y_{cm} = \frac{m_1y_1 + m_2y_2 + m_3y_3}{m_1 + m_2 + m_3} = \frac{3\sqrt{3}}{12}m$$

22. Two fly wheels are connected by a non-slipping belt as shown in the figure. $I_a = 4 \text{ kg m}^2$, $r_1 = 20$ cm, $I_2 = 20 \text{ kg m}^2$ and $r_2 = 30$ cm. A torque of 10 Nm is applied on the smaller wheel. Then match the entries of column I with appropriate entries of column II.

I	Quantities	II	Their numerical values (in SI units)
(a)	Angular acceleration of smaller wheel	(i)	$\frac{5}{3}$
(b)	Torque on the larger wheel	(ii)	$\frac{100}{3}$
(c)	Angular acceleration of larger wheel	(iii)	$\frac{5}{2}$



- 1) a-iii, b- ii, c-i 2) a-ii, b- iii, c-i 3) a- iii, b-i, c-ii 4) a- ii, b-i, c-iii **Ans: (1)**

Solution: $\tau_1 = I_1\alpha_1$

$$\alpha_1 = \frac{10}{4} = 2.5 \text{ rads}^{-2}$$

a is same for both

$$r_1\alpha_1 = r_2\alpha_2$$

$$\alpha_2 = \frac{r_1\alpha_1}{r_2} = \frac{2}{3} \times 2.5 = \frac{5}{3} \text{ rads}^{-2}$$

$$\tau_2 = I_2\alpha_2$$

$$= 20 \times \frac{5}{3} = \frac{100}{3} \text{ Nm}$$

23. If r_p , v_p , L_p and r_a , v_a , L_a are radii, velocities and angular momenta of a planet at perihelion and aphelion of its elliptical orbit around the Sun respectively, then

1) $r_p < r_a$, $v_p < v_a$, $L_p < L_a$

2) $r_p > r_a$, $v_p > v_a$, $L_p > L_a$

3) $r_p < r_a$, $v_p > v_a$, $L_p = L_a$

4) $r_p > r_a$, $v_p < v_a$, $L_p = L_a$

Ans: (3)

Solution: Conceptual

24. The total energy of a satellite in a circular orbit at a distance $(R + h)$ from the center of the Earth varies as $[R$ is the radius of the Earth and h is the height of the orbit from Earth's surface]

1) $\frac{1}{(R + h)}$

2) $-\frac{1}{(R + h)}$

3) $\frac{1}{(R + h)^2}$

4) $-\frac{1}{(R + h)^2}$

Ans. (2)

Solution: $TE = \frac{-GMm}{2(R + h)}$

25. Two wires A and B are made of same material. Their diameters are in the ratio of 1:2 and lengths are in the ratio of 1:3. If they are stretched by the same force, then increase in their lengths will be in the ratio of

(1) 4:3

(2) 3:4

3) 2 : 3

(4) 3:2

Ans. (1)

Solution: $Y = \frac{F\ell}{A\Delta\ell}$; $\Delta\ell \propto \frac{\ell^2}{r^2}$; $\frac{\Delta\ell_1}{\Delta\ell_2} = \frac{1}{3} \times \left(\frac{2}{1}\right)^2 = \frac{4}{3}$

26. A horizontal pipe carries water in a streamlined flow. At a point along the pipe, where the cross-sectional area is 10 cm^2 , the velocity of water is 1 ms^{-1} and the pressure is 2000 Pa . What is the pressure of water at another point where the cross-sectional area is 5 cm^2 ?

[Density of water = 1000 kgm^{-3}]

(1) 200 Pa

(2) 300 Pa

3) 400 Pa

(4) 500 Pa


Ans. (4)

Solution: $P_1 + \frac{1}{2}\rho V_1^2 = P_2 + \frac{1}{2}\rho V_2^2$

$2000 + \frac{1}{2} \times 1000 \times 1^2 = P_2 + \frac{1}{2} \times 1000 \times 4 \quad \therefore V_2 = \frac{A_1 V_1}{A_2}$

$P_2 = 500 \text{ Pa}$

27. Three metal rods of the same material and identical in all respects are joined as shown in the figure. The temperatures at the ends of these rods are maintained as indicated. Assuming no heat energy loss occurs through the curved surfaces of the rods, the temperature at the junction x is



1) 45°C

2) 60°C

3) 30°C

4) 20°C

Ans. (2)

Solution: $\theta - 0 + \theta - 90 + \theta - 90 = 0$

$3\theta = 180$

$\theta = \frac{180}{3} = 60^\circ\text{C}$

28. A gas is taken from state A to state B along two different paths 1 and 2. The heat absorbed and work done by the system along these two paths are Q_1 and Q_2 and W_1 and W_2 respectively. Then

(1) $Q_1 = Q_2$

(2) $W_1 = W_2$

(3) $Q_1 - W_1 = Q_2 - W_2$

(4) $Q_1 + W_1 = Q_2 + W_2$

Ans. (3)

Solution: $\Delta U_1 = \Delta U_2$

$Q_1 - W_1 = Q_2 - W_2$


29. At 27°C temperature, the mean kinetic energy of the atoms of an ideal gas is E_1 . If the temperature is increased to 327°C, then the mean kinetic energy of the atoms will be

- 1) $\frac{E_1}{2}$ 2) $\frac{E_1}{\sqrt{2}}$ 3) $\sqrt{2} E_1$ 4) $2E_1$ Ans. (4)

Solution: $E \propto T$

$$\frac{E_2}{E_1} = \frac{600}{300} = 2$$

30. The variations of kinetic energy $K(x)$, potential energy $U(x)$ and total energy as a function of displacement of a particle in SHM is as shown in the figure. The value of $|x_0|$ is



- 1) $\frac{A}{2}$ 2) $2A$ 3) $\frac{A}{\sqrt{2}}$ 4) $\sqrt{2}A$ Ans. (3)


Solution: Conceptual

31. The angle between the particle velocity and wave velocity in a transverse wave is [except when the particle passes through the mean position]

- (1) Zero radian (2) $\frac{\pi}{4}$ radian (3) $\frac{\pi}{2}$ radian 4) π radian Ans. (3)

Solution: Conceptual


32. A metallic sphere of radius R carrying a charge q is kept at certain distance from another metallic sphere of radius $R/4$ carrying a charge Q . What is the electric flux at any point inside the metallic sphere of radius R due to the sphere of radius $R/4$?



- 1) $\frac{Q}{\epsilon_0}$ 2) $\frac{Q}{\epsilon_0} - \frac{q}{\epsilon_0}$ 3) Zero 4) $\frac{q}{\epsilon_0} - \frac{Q}{\epsilon_0}$ Ans. (3)

Solution: According to gauss law

33. You are given a dipole of charge $+q$ and $-q$ separated by a distance $2R$. A sphere 'A' of radius ' R ' passes through the centre of the dipole as shown below and another sphere 'B' of radius ' $2R$ ' passes through the charge $+q$. Then the electric flux through the sphere A is



- 1) $-q/\epsilon_0$ 2) q/ϵ_0 3) Zero 4) $2q/\epsilon_0$ Ans. (1)


Solution: According to gauss law

34. A potential at a point A is -3 V and that at another point B is 5 V . What is the work done in carrying a charge of 5 C from B to A?

- (1) -40 J (2) -0.04 J (3) -0.4 J (4) -4 J **Ans. (2)**

Solution: $W = q[V_A - V_B] = 5 \times 10^{-3}[-3 - 5] = -0.04\text{ J}$

35. Charges are uniformly spread on the surface of a conducting sphere. The electric field from the centre of sphere to a point outside the sphere varies with distance r from the centre as



Solution: $E_{\text{in}} = 0$

$$E_{\text{out}} \propto \frac{1}{r}$$

36. Match Column-I with Column-II related to an electric dipole of dipole moment \vec{p} that is placed in a uniform electric field \vec{E} .

Column -I Angle between \vec{p} and \vec{E}	Column-II Potential energy of the dipole
a) 180°	(i) $-pE$
b) 120°	(ii) pE
c) 90°	(iii) $\frac{1}{2} pE$
	(iv) zero

- 1) a - ii, b - iii, c - iv 2) a-i, b - ii, c - iii 3) a - ii, b - iii, c - i 4) a-ii, b - i, c - iv **Ans. (1)**

Solution: $U = pE \cos \theta$

37. Which of the following statements is not true?

- 1) Electric field is always perpendicular to an equipotential surface.
 2) Work done to move a charge on an equipotential surface is not zero
 3) Equipotential surfaces are the surfaces where the potential is constant
 4) Equipotential surfaces for a uniform electric field are parallel and equidistant from each other

Ans. (2)


Solution: Conceptual

38. Which of the following is a correct statement?

- 1) Gauss's law is true for any closed surface
 2) Gauss's law is true for any open surface
 3) Gauss's law is not applicable when charges are not symmetrically distributed over a closed surface.
 4) Gauss's law does not hold good for a charge situated outside the Gaussian surface. **Ans. (1)**

Solution: Conceptual

39. In the following circuit, the terminal voltage across the cell is




- (1) 0.52 V (2) 1.68 V (3) 1.95 V (4) 2.71 V **Ans. (3)**

Solution: $I = \frac{\epsilon}{R + r} = 0.5\text{ A}$

$$V = \epsilon - Ir = 2 - 0.5 \times 0.1 = 1.95\text{ V}$$

40. Two cells of emfs E_1 and E_2 and internal resistances r_1 and r_2 ($E_2 > E_1$ and $r_2 > r_1$) respectively, are connected in parallel as shown in figure. The equivalent emf of the combination is E_{eq} . Then




- (1) $E_1 < E_{eq} < E_2$ and E_{eq} is nearer E_1 (2) $E_1 < E_{eq} < E_2$ and E_{eq} is nearer E_2
 (3) $E_{eq} > E_2$ (4) $E_{eq} < E_1$

Ans. (1)

Solution: $E_{eq} = \frac{E_1 r_2 + E_2 r_1}{r_1 + r_2}$

41. The variations of resistivity ρ with absolute temperature T for three different materials X, Y and Z are shown in the graph below. Identify the materials X, Y and Z.



- 1) X - copper, Y - nichrome, Z - semiconductor
 2) X - copper, Y - semiconductor, Z - nichrome
 3) X - semiconductor, Y - nichrome, Z - copper
 4) X-nichrome, Y - copper, Z – semiconductor

Ans. (1)

Solution: Based on nature


42. Given, a current carrying wire of non-uniform cross-section which of the following is constant throughout the length of wire?

- 1) Current, electric field and drift speed (2) Drift speed
 3) Current and drift speed (4) Current only

Ans. (4)

Solution: For a non-uniform cross-section only current through it is same.

43. The graph between variation of resistance of a metal wire as a function of its diameter keeping other parameters like length and temperature constant is



- 1) 2) 3) 4) **Ans. (4)**

Solution: $R \propto \frac{\ell}{D^2}$

44. Two thin long parallel wires separated by a distance 'r' from each other in vacuum carry a current of I ampere in opposite directions. Then, they will

- 1) Attract each other with a force per unit length of $\frac{\mu_0 I^2}{2\pi r^2}$
 (2) Attract each other with a force per unit length of $\frac{\mu_0 I^2}{2\pi r}$
 (3) Repel each other with a force per unit length of $\frac{\mu_0 I^2}{2\pi r}$
 (4) Repel each other with a force per unit length of $\frac{\mu_0 I^2}{2\pi r^2}$

Ans. (3)

Solution: Unlike current repel

$$\frac{F}{\ell} = \frac{\mu_0}{4\pi} \frac{2I^2}{r} = \frac{\mu_0 I^2}{2\pi r}$$

45. A solenoid is 1 m long and 4 cm in diameter. It has five layers of windings of 1000 turns each and carries a current of 7A. The magnetic field at the centre of the solenoid is
 1) 439.6T 2) $0.4396 \times 10^{-5}T$ (3) $4.396 \times 10^{-2}T$ (4) $43.96 \times 10^{-2}T$ **Ans. (3)**

Solution: $B = \mu_0 nI = 4\pi \times 10^{-7} \times 5 \times 10^3 \times 7 = 4.396 \times 10^{-2} T$

46. Two similar galvanometers are converted into an ammeter and a milliammeter. The shunt resistance of ammeter as compared to the shunt resistance of milliammeter will be
 (1) Equal 2) Zero (3) More (4) Less **Ans. (4)**

Solution: Conceptual


47. Which of the following statements is true in respect of diamagnetic substances?
 1) Susceptibility is small and negative 2) They are feebly attracted by magnets
 3) Permeability is greater than 1000. 4) Susceptibility decreases with temperature. **Ans. (1)**

Solution: Conceptual

48. Identify the correct statement
 (1) The magnetic field inside a solenoid is non uniform
 (2) A current carrying conductor produces an electric field around it.
 (3) A straight current carrying conductor has circular magnetic field lines around it.
 (4) The direction of magnetic field due to a current element is given by Flemings Left Hand Rule
Ans. (3)

Solution: Conceptual

49. Which of the following, graphs represents the variation of magnetic field B with perpendicular distance 'r' from an infinitely long, straight conductor carrying current?




Solution: $B \propto \frac{1}{r}$

50. If we consider an electron and a photon of same de-Broglie wavelength, then they will have same
 (1) Momentum 2) Angular momentum (3) Energy (4) Velocity **Ans. (1)**


Solution: $\lambda = \frac{h}{mv}$. As λ is constant, $p = mv$ is constant

51. The anode voltage of a photocell is kept fixed. The frequency of the light falling on the cathode is gradually increased. Then the correct graph which shows the variation of photo current I with the frequency ν of incident light is



Solution: Photo current is independent of frequency of incident radiation

52. When a bar magnet is pushed towards the coil, along its axis, as shown in the figure, the galvanometer pointer deflects towards X. When this magnet is pulled away from the coil, the galvanometer pointer



- (1) deflects towards X (2) deflects towards X'
(3) does not deflect (4) oscillates

Ans. (2)

Solution: Conceptual

53. A square loop of side 2 m lies in the Y-Z plane in a region having a magnetic field $\vec{B} = (5\hat{i} + 3\hat{j} - 4\hat{k})\text{T}$. The magnitude of magnetic flux through the square loop is

- (1) 10 Wb (2) 20 Wb (3) 12 Wb (4) 16 Wb **Ans. (2)**

Solution: $A = l \times b = 4\text{m}^2$

$$\phi = B_x A = 5 \times 4 = 20 \text{ Wb}$$

54. In domestic electric mains supply, the voltage and the current are

- (1) AC voltage and AC current (2) AC voltage and DC current
(3) DC voltage and DC current (4) DC voltage and AC current **Ans. (1)**

Solution: Conceptual

55. A sinusoidal voltage produced by an AC generator at any instant t is given by an equation $V = 311 \sin 314 t$. The rms value of voltage and frequency are respectively

- 1) 200 V, 100 Hz (2) 200 V, 50 Hz (3) 220 V, 100 Hz (4) 220 V, 50 Hz **Ans. (4)**

Solution: $\omega = 2\pi f = 314$, $f = 50 \text{ Hz}$, $V_{\text{rms}} = \frac{1}{\sqrt{2}} V_0 = \frac{1}{\sqrt{2}} \times 311$

56. A series LCR circuit containing an AC source of 100 V has an inductor and a capacitor of reactances 24Ω and 16Ω respectively. If a resistance of 6Ω is connected in series, then the potential difference across the series combination of inductor and capacitor only is

- 1) 40 V (2) 80 V (3) 400 V (4) 8 V **Ans. (2)**

Solution: $I = \frac{V}{Z} = \frac{100}{\sqrt{6^2 + 8^2}} = 10 \text{ A}$

$$V_L = IX_L = 10 \times 24 = 240 \text{ V}$$

$$V_C = IX_C = 10 \times 16 = 160 \text{ V}$$

$$V_L - V_C = 240 - 160 = 80 \text{ V}$$

57. Match the following types of waves with their wavelength ranges

	Waves		Wavelength ranges
(i)	Microwave	(a)	700 nm to 400 nm
(ii)	Visible light	(b)	1 nm to 10^{-3} nm
(iii)	Ultraviolet	(c)	0.1 m to 1 mm
(iv)	X-rays	(d)	400 nm to 1 nm

- (1) i-a, ii-d, iii -b, iv-c (2) i-c, ii-a, iii -d, iv-b
(3) i - d, ii-b, iii - c, iv - a (4) i - b, ii-c, iii-a, iv - d

Ans. (2)

Solution: Conceptual

58. A ray of light passes from vacuum into a medium of refractive index n . If the angle of incidence is twice the angle of refraction, then the angle of incidence in terms of refractive index is

- 1) $\cos^{-1}\left(\frac{n}{2}\right)$ 2) $\sin^{-1}\left(\frac{n}{2}\right)$ 3) $2\cos^{-1}\left(\frac{n}{2}\right)$ 4) $2\sin^{-1}\left(\frac{n}{2}\right)$ **Ans. (3)**


Solution: $i = 2r$; $n_1 \sin i = n_2 \sin r$

$$\sin i = n \sin \frac{i}{2}$$

$$2 \sin \frac{i}{2} \cos \frac{i}{2} = n \sin \frac{i}{2}$$

$$i = 2 \cos^{-1}\left(\frac{n}{2}\right)$$

59. A convex lens has power P . It is cut into two halves along its principal axis. Further one piece (out of two halves) is cut into two halves perpendicular to the principal axis as shown in figure. Choose the incorrect option for the reported lens pieces



- 1) Power of L_1 is $\frac{P}{2}$ 2) Power of L_2 is $\frac{P}{2}$ 3) Power of L_3 is $\frac{P}{2}$ 4) Power of L_1 is P **Ans. (1)**

Solution: L_1 no change

L_2 & L_3 have $\frac{P}{2}$

60. The image formed by an objective lens of a compound microscope is

- (1) Virtual and diminished (2) Real and diminished
(3) Real and enlarged (4) Virtual and enlarged

Ans. (3)

Solution: From image formation